

Reduced Environmental Stimulation Therapy (REST) in anxiety and depression: An experience sampling study[☆]



McKenna M. Garland^{a,b}, Raminta Wilson^a, Emily Adamic^{a,c}, Wesley K. Thompson^a, Armen C. Arevian^d, Murray B. Stein^{e,f,g}, Martin P. Paulus^{a,e,i}, Justin S. Feinstein^h, Sahib S. Khalsa^{a,i,*}

^a Laureate Institute for Brain Research (LIBR), University of Tulsa, Tulsa, OK, USA

^b Kendall College of Arts and Sciences, University of Tulsa, Tulsa, OK, USA

^c College of Engineering and Natural Sciences, University of Tulsa, Tulsa, OK, USA

^d Chorus Innovations, LLC, Los Angeles, CA, USA

^e Department of Psychiatry, University of California San Diego, La Jolla, CA, USA

^f Herbert Wertheim School of Public Health, University of California San Diego, La Jolla, CA, USA

^g Psychiatry Service, Veterans Affairs San Diego Healthcare System, San Diego, CA, USA

^h Float Research Collective, Kihai, HI, USA

ⁱ Oxley College of Health Sciences, University of Tulsa, Tulsa, OK, USA

ARTICLE INFO

Keywords:

Reduced Environmental Stimulation Therapy
Floatation-REST
Float therapy
Anxiety disorder
Major depressive disorder
Ecological momentary assessment

ABSTRACT

Background: Reduced Environmental Stimulation Therapy (REST) is a behavioral intervention that systematically attenuates external sensory input to the nervous system. Previous studies have demonstrated acute anxiolytic and antidepressant effects of single sessions of REST in anxious individuals, however the duration and time course of these effects is unknown. In the current study, we used experience sampling and multiple sessions of REST to explore the time course of the anxiolytic and antidepressant effects over a 48-hour time period.

Methods: 75 adults with anxiety and/or depression were randomized to complete 6 sessions of REST (either pool-REST, pool-REST preferred, or chair-REST). Post-REST effects were tracked using experience sampling on a smartphone with a ten-item survey administered at 6 time points for each session (i.e., immediately before and after REST and then 4, 8, 24, and 48 h later). Using principal component analysis, responses to survey items at baseline were reduced to broad symptom clusters of anxiety, depression, and serenity and were utilized in linear mixed effects models to determine the magnitude and time course of post-REST effects.

Results: REST was associated with significant decreases in anxiety and depression, and significant increases in serenity, with effects lasting for 48 h. Repeated exposure to REST was associated with lower baseline levels of anxiety at later sessions.

Conclusion: These initial findings suggest that the anxiolytic and antidepressant effects of REST persist for at least two days, and that repeated REST sessions may have additive effects on lowering anxiety-related symptoms. These findings could help to determine the optimal intervention frequency of REST and facilitate future investigations focused on the combination of REST with standard treatments for anxiety and depression.

Introduction

Reduced Environmental Stimulation Therapy (REST) is a poorly understood non-pharmacological stress-reduction intervention designed to systematically attenuate external sensory input to the nervous system [1]. Typically, REST involves floating effortlessly in a shallow

pool of warm water that is saturated with Epsom salt (i.e., ‘pool-REST’). Pool-REST environments are lightproof, soundproof, and humidity and temperature-controlled, such that signals from visual, auditory, olfactory, gustatory, thermosensory, tactile, vestibular, and proprioceptive channels are minimized, as is movement and speech. ‘Chair-REST’ is another form of REST that involves reclining in an ergonomically-

[☆] Clinical Trial Registration Identifier; NCT03899090

* Corresponding author at: Laureate Institute for Brain Research (LIBR), University of Tulsa, Tulsa, OK, USA.

E-mail address: skhalsa@laureateinstitute.org (S.S. Khalsa).

engineered “zero-gravity” chair designed to take pressure off the spinal cord; the chair is placed in a room with dim lighting and quiet surroundings, similar but not identical to what would be experienced during pool-REST, thus serving as an active comparator.

Perhaps the most consistent clinical observation to date is that REST induces acute anxiolytic and antidepressant symptom reductions [2–5]. However, the precise duration and time course of these effects are unknown. Here, we explored the time course of acute REST effects using experience sampling over a 48-hour time period after six sessions, with additional exploratory analyses to understand the additive influence of repeated REST sessions on anxiety and depression symptoms.

Method

Participants

75 treatment-seeking adults with anxiety and/or depression, and elevated levels of anxiety (as measured by an OASIS score ≥ 6) and anxiety sensitivity (as measured by an ASI-3 total score ≥ 24) were recruited through LIBR’s participant databases and from the Tulsa community. The full protocol and eligibility criteria are described elsewhere [6]. The study was pre-registered (clinicaltrials.gov NCT03899090). All study procedures were approved by the Western IRB. All participants provided written informed consent before participation and received compensation.

Procedures

75 participants were randomized to 6 sessions of pool-REST ($n = 25$), pool-REST preferred ($n = 25$), or chair-REST ($n = 25$). The time course of REST effects was assessed at 6 time points: before and after each REST session via iPad in the laboratory and then 4, 8, 24, and 48 h later via text-message links sent to the participant’s cellular phone using the web-based Chorus platform (Chorus Inc., Los Angeles CA).

The survey consisted of a ten-item questionnaire assessing symptoms known to be influenced by REST environments [3], including the PANAS-X Serenity subscale [7], Karolinska Sleepiness Scale [8], and continuous Visual Analogue Scales (VAS) assessing well-being, stress/anxiety, depression, peacefulness, relaxation, and degree of muscle tension. All survey item values were converted into z-scores before analysis.

Experimental Conditions

Pool-REST

This involved 1-hour session durations prescribed at fixed 1-week intervals within an open or enclosed circular float pool with an 8-foot diameter (Floataway Inc., Norfolk, UK).

Pool-REST Preferred

This involved up to 2-hour session durations. Participants were allowed flexibility in arranging their floating sessions to match their preferred schedule within a 12-week period, with the only requirement being that there needed to be a minimum of 24 h between sessions.

Chair-REST

This involved reclining in a Zero Gravity Chair (PC510, Classic Power, Series 2, Human Touch Inc., Long Beach, CA). This active comparator closely matched the pool-REST intervention on many parameters including a supine body position in a dimly lit and quiet room, a 1-hour session duration prescribed at fixed 1-week intervals, and a similar instruction set emphasizing the importance of stillness and wakefulness throughout each session.

Statistical Analysis

All analyses were performed in RStudio 3.6.0. Using a principal component (PC) analysis, responses to survey items at the initial pre-float session time point were reduced to broad symptom clusters using the Kaiser-Guttman criterion and visual inspection of scree plots. A promax rotation was applied to the retained PCs. Survey items that loaded most strongly onto each PC were assigned exclusively to that scale. Once PC scales were generated, individual item responses were converted to POMP scores (standardized units representing the “Percent Of Maximum Possible,” ranging from 0% to 100%). Unit-weighted PC scales were then created by summing POMP scores for each respective component, and this was replicated for each of the subsequent survey timepoints. The resulting unit-weighted PC scales were utilized in independent linear mixed effects (LME) models to determine the magnitude and time course of REST effects. A hypothesis-wise Bonferroni adjustment across the variables of interest was performed, whereby alpha was divided by the number of PCs extracted ($\alpha = 0.05/3 = 0.017$). LME modeling allowed for the examination of main effects of REST session, survey timepoint, and condition, and the interaction between these variables (i.e., $PC\# \sim Survey*Session*Condition$). The models included subject ID and REST session as random effects (formula: $\sim 1|Subject\ ID/Session$) and utilized an AR1 covariance structure. *Post hoc* two-sided t-tests with Holm corrections were used to interpret significant main effects and simple effects for all significant interactions.

Results

Participants

75 participants (mean age = 35.0, SD = 11.3, 77% female, 81% non-Hispanic White) were randomized. All participants had a DSM-5 diagnosis of an anxiety, stress-related, or depressive disorder, with the most common psychiatric diagnoses being major depressive disorder (97.3%) and generalized anxiety disorder (50.7%). At baseline, participants reported elevated anxiety and depression symptoms (average OASIS = 9.7; average PHQ-9 = 12.1, respectively) and high anxiety sensitivity (average ASI-3 = 40.4). The groups did not differ significantly on any sociodemographic or clinical variables at baseline [6]. Of the 25 participants randomized to each group, the six session completer numbers were: 17 (chair-REST), 19 (pool-REST), and 21 (pool-REST preferred); see [6] for further details.

Principal component analysis

The PC analysis suggested a three-component solution, with the top PCs (Eigenvalues: 4.43, 1.76, and 1.11) cumulatively accounting for 74% of the total variance. The top PCs in descending order of eigenvalue were designated: Serenity, Anxiety, and Depression, and accounted for 45%, 18%, and 11% of the total variance, respectively.

Magnitude and duration of REST effects

Anxiety

In the LME evaluating self-reported anxiety, a significant main effect of survey ($F(5) = 94.30, p < 0.001, \eta_p^2 = 0.25$) and REST session ($F(5) = 5.10, p = 0.0002, \eta_p^2 = 0.09$), and a significant interaction between survey and condition ($F(10) = 2.88, p = 0.0015, \eta_p^2 = 0.02$) was observed. *Post hoc* comparisons revealed reductions in self-reported anxiety lasting 48 h across all three REST interventions ($ps \leq 0.01$, Cohen d s (d) = 0.47–1.64; Fig. 1). Additive effects on anxiety levels were observed with significantly lower anxiety at sessions 5 ($p = 0.0453, d = 0.30$) and 6 ($p = 0.0112, d = 0.35$) versus session 1 (Fig. 1).




Fig. 1. Top: Time course of average REST effects across six sessions. Significant reductions in anxiety lasted for 48 h across all three REST conditions. Significant decreases in depression were present for 48 h for the pool-REST preferred condition and up to 8 h for the chair-REST and pool-REST conditions. Significant increases in serenity ratings were present at all time points for both pool-REST conditions but were no longer present at 48 h for the chair-REST condition. **Bottom:** Additive effects of REST. Pre-session anxiety at visits 5 and 6 was significantly lower than pre-session anxiety ratings at visit 1. There were no significant additive effects for depression or serenity. Ribbons reflect the standard error of the mean. PC: Principal Component. REST: Reduced Environmental Stimulation Therapy.

Depression

In the LME evaluating self-reported depression, a significant main effect of survey ($F(5) = 87.96, p < 0.0001, \eta_p^2 = 0.23$), and a significant interaction between survey and condition ($F(10) = 2.29, p = 0.0116, \eta_p^2 = 0.02$) was observed. *Post-hoc* comparisons revealed significant reductions in self-reported depression lasting 48 h for the pool-REST preferred intervention ($ps < 0.001, ds = 0.52-1.64$) but only up to 8 h in the chair-REST ($p = 0.0001, d = 0.65$) and pool-REST interventions ($p = 0.0002, d = 0.57$; Fig. 1).

Serenity

In the LME evaluating self-reported serenity, a significant main effect of survey ($F(5) = 198.62, p < 0.0001, \eta_p^2 = 0.41$) and a significant interaction between survey and condition ($F(10) = 2.50, p = 0.006, \eta_p^2 = 0.02$) was observed. *Post hoc* comparisons revealed significantly higher levels of serenity at 24 h for the chair-REST intervention ($ps \leq 0.01; ds = 0.50-1.71$) and 48 h across the pool-REST and pool-REST preferred interventions ($ps \leq 0.01; ds = 0.38-2.38$; Fig. 1).

Discussion

This exploratory study examined the time course of the acute effects of REST in anxious and depressed adults. Significant reductions in anxiety and depression and increases in serenity were observed, lasting up to 48 h post-session. REST also had cumulative effects whereby lower anxiety symptoms were reported prior to the fifth and sixth sessions. These preliminary findings indicate that REST can provide persistent short-term affective symptom relief, suggesting its potential as a viable non-pharmacological intervention to be evaluated further.

Our results inform decisions regarding the optimal number and duration of REST sessions for future randomized clinical efficacy trials. Specifically, a 48-hour window captured most of the acute changes in anxious and depressed mood ratings. Setting a minimum 48-hour interval between REST sessions would thus seem feasible for a study

attempting to induce a more sustained form of anxiolysis. While the acute reductions in anxiety and depression were followed by some increases, the overall symptom reductions remained below baseline levels at the 48-hour timepoint. Thus, the upper limit of this clinical window remains uncharacterized. The additive anxiolytic effects observed at sessions five and six indicate that future REST studies should evaluate the impact of a larger number of sessions. It is currently unclear whether such additive effects would continue to grow with additional sessions, although this may be why prior REST studies have employed larger session numbers (e.g., ranging from 9 to 33 sessions) [4,5].

Several limitations must be acknowledged. The small sample size was not adequately powered for a between-group analysis of pool- vs. chair-REST (i.e., active comparator) effects in this open label exploratory study, nor was a placebo or sham intervention arm included. It is presently difficult to pinpoint the mechanism of the persistent effects of REST despite previously observed impacts on peripheral and central autonomic targets [9,10]. Although the survey items used in this study were extracted from standardized scales with acceptable psychometric properties [7,8] for the experience sampling approach, they do not replace clinical rating scales. Finally, the longer-term clinical effects of repeated sessions of REST remain uncharacterized in anxious and depressed individuals.

Ethics Statement

The study was approved by the Western Institutional Review Board under Protocol #20150528 and was performed in accordance with Declaration of Helsinki. All participants gave their written informed consent before participation and received compensation.

Funding

This research was supported by National Center for Complimentary and Integrative Health (NCCIH) R34AT009889. The authors are also

supported by the National Institute of Mental Health (R01MH127225, K23MH112949 to SSK), National Institute of General Medical Sciences Center Grant Award (1P20GM121312 to MPP, SSK), and The William K. Warren Foundation. The funding sources had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; or decision to submit the manuscript for publication.

Data Availability

Requests for sharing of data used in this analysis can be made to the corresponding author. Any sharing of data will be subject to obtaining appropriate agreements from the principal investigators or data custodians.

Declaration of Competing Interest

The authors declare the following financial interests/leadership roles which may be considered as potential competing interests: Armen C. Arevian (ACA) is founder of Insight Health Systems, Arevian Technologies, and Open Science Initiative. ACA developed the Chorus platform, which is licensed from the University of California Los Angeles to Insight Health Systems. Martin Paulus is an advisor to Spring Care, Inc., a behavioral health startup, and has received royalties for an article about methamphetamine in UpToDate. Justin Feinstein is the president and director of the nonprofit Float Research Collective. Sahib Khalsa is an executive committee member of the International Society of Contemplative Research and a board member of the Float Research Collective, both of which are unpaid roles. There are no other competing interests to report for any authors.

Acknowledgments

We thank David Watson, Henry Yeh, and Joel Greenshields for coding, statistical, and psychometrics consultations. We would also like

to acknowledge Laura Garrison for her contributions to data collection. Finally, we would like to thank the custodial staff at the Laureate Institute for Brain Research for their diligence in maintaining the REST facility. These results were initially presented at the Anxiety and Depression Association of America Conference, Denver, Colorado, in March 2022.

References

- 1 Feinstein JS, Khalsa SS, Yeh HW, Wohlrab C, Simmons WK, Stein MB, et al. Examining the short-term anxiolytic and antidepressant effect of Floatation-REST. *PLoS One* 2018;13(2):e0190292 <https://doi.org/10.1371/journal.pone.0190292>
- 2 Khalsa SS, Moseman SE, Yeh HW, Upshaw V, Persac B, Breeze E, et al. Reduced environmental stimulation in anorexia nervosa: an early-phase clinical trial. *Front Psychol* 2020;2534. <https://doi.org/10.3389/fpsyg.2020.567499>
- 3 Feinstein JS, Khalsa SS, Yeh H, Al Zoubi O, Arevian AC, Wohlrab C, et al. The elicitation of relaxation and interoceptive awareness using floatation therapy in individuals with high anxiety sensitivity. *Biol Psychiatry: Cogn Neurosci Neuroimaging* 2018;3(6):555–62. <https://doi.org/10.1016/j.bpsc.2018.02.005>
- 4 Bood SÅ, Sundequist U, Kjellgren A, Nordström G, Norlander T. Effects of floatation rest (restricted environmental stimulation technique) on stress related muscle pain: are 33 floatation sessions more effective than 12 sessions? *Soc Behav Pers: Int J* 2007;35(2):143–56. <https://doi.org/10.2224/sbp.2007.35.2.143>
- 5 Jonsson K, Kjellgren A. Promising effects of treatment with floatation-REST (restricted environmental stimulation technique) as an intervention for generalized anxiety disorder (GAD): a randomized controlled pilot trial. *BMC Complement Altern Med* 2016;16:1–12. <https://doi.org/10.1186/s12906-016-1089-x>
- 6 Garland MM, Wilson R, Thompson WK, Arevian A, Stein MB, Paulus MP, et al. A randomized controlled safety and feasibility trial of Floatation-REST in a sample of clinically anxious and depressed individuals. *MedRxiv* 2023. <https://doi.org/10.1101/2023.05.27.23290633>
- 7 Watson, D., & Clark, L.A. (1994). The PANAS-X: Manual for the positive and negative affect schedule-expanded form.
- 8 Åkerstedt T, Gillberg M. Subjective and objective sleepiness in the active individual. *Int J Neurosci* 1990;52(1–2):29–37. <https://doi.org/10.3109/00207459008994241>
- 9 Flux MC, Fine TH, Poplin T, Al Zoubi O, Schoenhals WA, Schettler J, et al. Exploring the acute cardiovascular effects of Floatation-REST. *Front Neurosci* 2022;16:995594 <https://doi.org/10.3389/fnins.2022.995594>
- 10 Al Zoubi O, Misaki M, Bodurka J, Kuplicki R, Wohlrab C, Schoenhals WA, et al. Taking the body off the mind: Decreased functional connectivity between somato-motor and default-mode networks following Floatation-REST. *Hum Brain Mapp* 2021;42(10):3216–27. <https://doi.org/10.1002/hbm.25429>

Unpublished Supplementary Materials

	Item	Original Scale
1.	In general, over the past few hours, how have you felt?	-10 (Pretty Bad) to 10 (Pretty Good) ^{a,c}
2.	Over the past few hours, how stressed or anxious have you felt?	0 (Not at All) to 100 (Extremely) ^c
3.	Over the past few hours, how sad, down, or depressed have you felt?	0 (Not at All) to 100 (Extremely) ^c
4.	Over the past few hours, how calm have you felt?	1 (Very Slightly or Not at All) to 5 (Extremely) ^b
5.	Over the past few hours, how relaxed have you felt?	1 (Very Slightly or Not at All) to 5 (Extremely) ^b
6.	Over the past few hours, how at ease have you felt?	1 (Very Slightly or Not at All) to 5 (Extremely) ^b
7.	How have you felt over the past few hours?	1 (Extremely Alert) to 9 (Extremely Sleepy) ^b
8.	Over the past few hours, how content or peaceful have you felt?	0 (Not at All) to 100 (Extremely) ^c
9.	Over the past few hours, how relaxed have you felt?	0 (Not at All) to 100 (Extremely) ^c
10.	Over the past few hours, how much muscle tension or tightness have you felt in your body?	0 (Not at All) to 100 (Extremely) ^c
*11.	How much of the effects from the float experience are you still feeling right now?	0% to 100% ^c
*12.	Please describe any effects that you are still feeling from the float experience.	Free response text box

Supplemental Table 1. *Experience Sampling Method Survey Items.* * Indicates items that were only assessed at post-float time periods. ^aItem was reverse coded prior to analyses. ^bItem used a Likert scale. ^cItem used a VAS scale.

Unpublished Supplementary Materials

REST Condition	Missing (n)	Expected (n)	Percentage Missing
Chair	113	666	17%
Pool	105	774	14%
Pool preferred	87	798	11%
TOTAL	305	2,238	14%

Supplemental Table 2. Missing Survey Data. Percentage of missing data split by REST condition. Expected values include only participants who remained in the study at the respective survey time point (i.e., does not include withdrawn participants). Missing data was observed for 14% of our primary experience sampling measures which is in line with (or slightly lower than) typical rates which hover around 21% (Wrzus & Neubauer, 2022).

Unpublished Supplementary Materials

Serenity (PC1) $\lambda = 4.43$	Anxiety (PC2) $\lambda = 1.76$	Depression (PC3) $\lambda = 1.11$
At ease* .43	Muscle tension .78	Sad/Down/Depressed .90
Peaceful .42	Stress/Anxiety .44	Recent well-being .34
Relaxed .42	Alertness .25	
Relaxed* .42		
Calm* .42		


Supplemental Table 3. Promax-rotated item loadings. The top three PCA factors in descending order of eigenvalue were designated as follows: Serenity, Anxiety, and Depression, and accounted for 45%, 18%, and 11% of the total variance, respectively. *Indicates items on the Serenity subscale of the PANAS-X.

Unpublished Supplementary Materials

	<i>Serenity Comparisons</i>									
	<i>Pre vs. Post</i>		<i>Pre vs. 4hr</i>		<i>Pre vs. 8hr</i>		<i>Pre vs. 24hr</i>		<i>Pre vs. 48hr</i>	
<i>Condition</i>	<i>p</i>	<i>d</i>	<i>p</i>	<i>d</i>	<i>p</i>	<i>d</i>	<i>p</i>	<i>d</i>	<i>p</i>	<i>d</i>
Chair-REST	< 0.01	1.71	< 0.01	0.92	< 0.01	0.88	< 0.01	0.50	0.08	0.33
Pool-REST	< 0.01	1.83	< 0.01	1.08	< 0.01	0.92	< 0.01	0.54	0.02	0.38
Pool-REST Preferred	< 0.01	2.38	< 0.01	1.54	< 0.01	1.46	< 0.01	0.83	< 0.01	0.58
	<i>Anxiety Comparisons</i>									
	<i>Pre vs. Post</i>		<i>Pre vs. 4hr</i>		<i>Pre vs. 8hr</i>		<i>Pre vs. 24hr</i>		<i>Pre vs. 48hr</i>	
<i>Condition</i>	<i>p</i>	<i>d</i>	<i>p</i>	<i>d</i>	<i>p</i>	<i>d</i>	<i>p</i>	<i>d</i>	<i>p</i>	<i>d</i>
Chair-REST	< 0.01	0.99	< 0.01	1.18	< 0.01	0.92	< 0.01	0.69	0.01	0.47
Pool-REST	< 0.01	1.42	< 0.01	1.03	< 0.01	0.85	< 0.01	0.61	< 0.01	0.63
Pool-REST Preferred	< 0.01	1.58	< 0.01	1.64	< 0.01	1.13	< 0.01	1.01	< 0.01	0.84
	<i>Depression Comparisons</i>									
	<i>Pre vs. Post</i>		<i>Pre vs. 4hr</i>		<i>Pre vs. 8hr</i>		<i>Pre vs. 24hr</i>		<i>Pre vs. 48hr</i>	
<i>Condition</i>	<i>p</i>	<i>d</i>	<i>p</i>	<i>d</i>	<i>p</i>	<i>d</i>	<i>p</i>	<i>d</i>	<i>p</i>	<i>d</i>
Chair-REST	< 0.01	1.03	< 0.01	0.76	< 0.01	0.65	0.10	0.35	0.79	0.12
Pool-REST	< 0.01	1.15	< 0.01	0.64	< 0.01	0.57	0.11	0.34	0.12	0.33
Pool-REST Preferred	< 0.01	1.64	< 0.01	1.20	< 0.01	1.02	< 0.01	0.80	< 0.01	0.52


Supplemental Table 4. Survey by Group Interaction Post-hoc Comparisons. *p* indicates Holm-corrected p-value associated with post-hoc comparison test of simple effects. *d* represents associated Cohen's D effect size.

Unpublished Supplementary Materials




Supplemental Figure 1. CONSORT Diagram. C = Completed. W = Withdrew.

Unpublished Supplementary Materials



Supplemental Figure 2. Average change score from pre-session ratings across six REST sessions split by condition.

Unpublished Supplementary Materials




Supplemental Figure 3. Residual Global REST Effects based on Item 11 of the survey (see Supplemental Table 1).

Unpublished Supplementary Materials




Supplemental Figure 4. Common Themes of Post-REST Effects based on Item 12 of the survey (see Supplemental Table 1). Colors are for illustrative purposes only. Word size is scaled to the relative frequency of each term.

Unpublished Supplementary Materials



Supplemental Figure 5. Change Scores Illustrating the Time Course of Effects Across All Six REST Sessions in Individuals Utilizing Psychotherapy ($n=19$) Versus Individuals Not Utilizing Psychotherapy ($n=56$). Psychotherapy utilization was measured at the pre-intervention visit. Ribbons indicate 95% confidence intervals around the mean.

Unpublished Supplementary Materials



Supplemental Figure 6. Change Scores Illustrating the Time Course of Effects Across All Six REST Sessions in Individuals Taking Psychotropic Medications ($n=28$) Versus Individuals Not Taking Psychotropic Medications ($n=47$). Psychotropic medications were measured at the pre-intervention visit. Ribbons indicate 95% confidence intervals around the mean.